

Oak Leaf-tier and Oak Leaf-roller



1. Bell-shaped appearance of adult oak leaf-tier



2. Oak leaf-tier larva on red oak bud



3. Oak leaf-tier damaged scarlet oak leaves



4. Oak leaf-roller damaged red oak leaves

The oak leaf-tier, *Croesia semipurpurana* (Kearfott), and the oak leaf-roller, *Archips semiferanus* (Walker), are important Tortricidae moths in a complex of native species feeding in the early spring on oak foliage throughout the forests of Eastern North America. Outbreaks periodically develop with frequent and severe defoliation resulting in tree mortality. The last outbreak during the 1960's and 1970's covered large areas of Connecticut, Massachusetts, New York, Pennsylvania, Virginia, and West Virginia.

Either moth species may dominate during an outbreak depending upon which oak species are present. Oak leaf-tier caterpillars tend to favor red, scarlet, black, and pin oaks; whereas, oak leaf-roller caterpillars feed not only on red and scarlet oaks, but also on chestnut and white oaks commonly found along ridgetops.

Resting adults of both species hold their wings in a characteristic bell shape. The adult wingspan of the oak leaf-tier is smaller (0.5 inch) than the oak leaf-roller (0.9 inch). Adults of both species are distinctly marked. The oak leaf-tier is yellow with a large, violet to brown front wing; the oak leaf-roller is creamy brown and gray with a darker oblique band across its front wing.

Despite their small size, larvae of both species, if abundant, can severely defoliate an oak tree. Newly hatched oak leaf-tier larvae enter opening buds and eat immature leaves. This feeding habit can destroy most new growth. Surviving leaves are deformed and riddled with holes. Oak leaf-roller larvae feed from leaves webbed together or rolled at the edges into shelters.

Larvae of both species exhibit various colors and markings as they mature. Immature larvae are several shades of green with a dark head capsule and thoracic shield and cannot easily be identified to species until they are mature. Mature oak leaf-tier larvae (left-facing) are 0.5 inch long with both a dark eye patch and a bar on the side of the head, and dark legs. Mature oak leaf-roller larvae (right-facing) are 1.2 inch long with either a dark eye patch or a dark bar on the head, and pale legs.



5. Mature, oak leaf-tier larva



6. Mature, oak leaf-roller larva

Photo Credits: Photo 3, Pennsylvania Bureau of Forestry
Photos 1-2 and 4-6, USDA Forest Service



For additional information,
contact:

USDA Forest Service
Forest Health Protection
180 Canfield Street
Morgantown, WV 26505
(304) 285-1541

USDA Forest Service
Forest Health Protection Unit
200 Weaver Boulevard
Asheville, NC 28802
(828) 257-4320